2014

Measuring the impact of contraceptive use on unintended pregnancy and other health outcomes

STEP UP Research Programme Consortium

Follow this and additional works at: https://knowledgecommons.popcouncil.org/departments_sbsr-rh

Part of the [Demography, Population, and Ecology Commons](#), [Family, Life Course, and Society Commons](#), [International Public Health Commons](#), [Maternal and Child Health Commons](#), [Obstetrics and Gynecology Commons](#), and the [Women's Health Commons](#)

How does access to this work benefit you? Let us know!

Recommended Citation

"Measuring the impact of contraceptive use on unintended pregnancy and other health outcomes," STEP UP Brief. New York: Population Council.

This Brief is brought to you for free and open access by the Population Council.

MEASURING THE IMPACT OF CONTRACEPTIVE USE ON UNINTENDED PREGNANCY AND OTHER HEALTH OUTCOMES

National and global support for services that enable women and their partners to have access to a full range of quality contraceptives are growing rapidly, as evidenced through the financial commitments made during the London Summit on Family Planning, and the ensuing FP2020 Initiative that seeks to reduce unmet need for such services by 120 million women.

With these investments, and the anticipated increases in effective contraceptive use, come expectations of substantial and demonstrable impacts on a range of health outcomes, including significant reductions in unintended pregnancies, maternal, newborn and child deaths and unsafe abortions.

Attributing health outcomes to a specific health intervention is always challenging, and direct measurement of outcomes averted is not possible. Consequently, several statistical models have been developed that estimate the number of health outcomes that will be averted because of contraceptive use. The table on page 3 describes and compares five of the most commonly used approaches: *Adding it Up* (AIU); *Impact2*; *ImpactNow*; *Reality Check*; and *FamPlan/LiST*. AIU is used by staff at the Guttmacher Institute for undertaking and presenting analyses of outcomes and is not in the public domain; the other four estimation models have been developed for use at national and sub-national levels by donors, programme managers and policymakers. While these are the most frequently used approaches to measure the impact of contraceptive use, other models are available that address similar

APPROACHES FOR MEASURING THE IMPACT OF CONTRACEPTIVE USE

Adding it up (AIU)

ORGANISATION: Guttmacher Institute
 CONTACT: Jacqueline E. Darroch
 (JEDarroch@guttmacher.org)
 WEBSITE: Darroch, J, Singh S. 2011. *Adding It Up: The Costs And Benefits Of Investing In Family Planning And Maternal And Newborn Health—Estimation Methodology*. New York: Guttmacher Institute.
www.guttmacher.org/pubs/AIU-methodology.pdf

Impact2

ORGANISATION: Marie Stopes International
 CONTACT: Michelle Weinberger
 (michelle.weinberger@mariestopes.org)
 WEBSITE: Marie Stopes International. 2012. *Impact 2: An innovative tool for measuring the impact of reproductive health programmes*. London: Marie Stopes International.
www.mariestopes.org/impact-2

ImpactNow

ORGANISATION: USAID Health Policy Project, Futures Group
 CONTACT: Ellen Smith
 (esmith@futuresgroup.com)
 WEBSITE: Not yet available

Reality Check

ORGANISATION: USAID RESPOND Project, EngenderHealth
 CONTACT: Melanie Yahner
 (myahner@engenderhealth.org)
 WEBSITE: Not yet available www.respond-project.org

FamPlan/LiST

ORGANISATION: USAID Health Policy Project, Futures Institute
 CONTACT: John Stover
 (jstover@futuresinstitute.org)
 WEBSITE: Spectrum Manual: Spectrum System of Policy Models: www.futuresinstitute.org/spectrum.aspx

questions. The *Crosswalk* document¹ helps advocates, programme planners, decision makers, and others select which among these tools might best address their goals or questions, as well as provide insight into how to interpret the outputs of each tool.

However, concerns have been expressed that this range of approaches—which overlap in the impacts they estimate, but sometimes use different inputs, mathematical algorithms and assumptions, and may produce different results—may be confusing for policy makers, managers and donors who want to measure or evaluate these impacts. To address these concerns, the STEP UP consortium convened two expert meetings (in September 2013 and March 2014) to review the estimation methods used; come to consensus on methodologies and assumptions, where possible; and provide guidance to those wanting to measure such impacts. Following the first meeting, the same data from four countries (Bangladesh, Burkina Faso, Colombia, Malawi) were inputted into each model and the estimates generated for numerous impacts were compared. Substantial differences emerged, which provided the basis for intensive discussions between the model developers during the second meeting. These discussions led to modifications in the models through reaching a consensus on the following core elements.

PREGNANCY RATES

Pregnancy rates are important for determining the likelihood of a pregnancy among sexually active, fecund women who do not want to become pregnant when contraception is not used. Agreement was reached that the same pregnancy rate for a “non-user at risk of unintended pregnancy” should be used as the default where applicable in all models and kept constant for the foreseeable future. Using the AIU methodology, the estimated pregnancy rate among women with unmet need using no method across 148 developing countries is 31%, with an inter-quartile plausibility range of 23–38%. This rate is calculated by dividing the number of unintended pregnancies among the estimated number of women with unmet need using no method, by the total number of women with unmet need using no method. This rate is lower than some previous estimates. It should be noted that this default pregnancy rate will be adopted by the models *Impact2*, *ImpactNow*, *Reality Check* and *FamPlan* but the published tables in *Adding It Up* will continue to use Guttmacher’s estimates of regional pregnancy rates.

CONTRACEPTIVE METHOD FAILURE RATES

Agreement was reached that the same method failure rates should be used where applicable in all models because it is important to ensure that the estimated impact of using contraception takes into account the fact that no method is 100% effective. The group recommended using the method-specific rates calculated by Cleland, Ali & Shah¹ and by Trussell.² Moreover, agreement was reached that these rates should not be adjusted for individual countries because: i) it is not possible to distinguish unintended pregnancy by method used in national settings; and ii) these global estimates are widely accepted. As with the pregnancy rates, this consensus will be implemented by the models *Impact2*, *ImpactNow*, *Reality Check* and *FamPlan*; the *Adding It Up* analysis will continue to be based on the adjusted failure rates estimated by Guttmacher staff, which differ by region.

ESTIMATING ABORTIONS AND MISCARRIAGES AVERTED

AIU, *Reality Check*, and *FamPlan* will use the “% of unintended pregnancies ending in abortion” rates estimated by the Guttmacher Institute. *Impact2* and *ImpactNow* currently apply sub-regional abortion ratios to the number of live births averted, and will explore using the Guttmacher Institute’s unintended pregnancy ratios instead. The models estimate the number of miscarriages averted slightly differently, but the differences are very minor and so no change was deemed necessary.

MATERNAL DEATHS AVERTED

Published national maternal mortality ratios (MMR) represent the overall risk of maternal mortality given the national distribution of pregnancy outcomes (births, abortions, miscarriages). Applying these ratios to estimates of live births averted by contraceptive use does not accurately represent the risk of maternal mortality of an adjusted distribution on unintended pregnancies (see above). *Reality Check* and *FamPlan* will adapt the *AIU* approach to adjusting the national maternal mortality ratio estimations,⁴ to reflect the adjusted distribution of unintended pregnancy outcomes. This adjustment will be done using WHO sub-regional unsafe abortion rates and a Guttmacher Institute estimate of two deaths per 100,000 safe abortions. *Impact2* and *ImpactNow* will explore using this adjusted approach as part of exploring adjusting pregnancy outcomes based on pregnancy intention.

1 Godbole R, Smith E. 2012. *Crosswalk of Family Planning Tools: A Guide to Costing, Planning, and Impact Analysis Tools*. Washington, DC: Health Policy Project, USAID, Bureau for Global Health.

2 Cleland J, Ali MM, and Shah I. 2006. “Dynamics of contraceptive use”, in: UN Department of Economic and Social Affairs, Population Division, *Levels and Trends of Contraceptive Use as Assessed in 2002*, pp. 87–115, Table 22, page 98.

3 Trussell J. 2007. “Contraceptive efficacy,” in: Hatcher RA et al. (eds.), *Contraceptive Technology*, 19th ed., New York: Ardent Media.

4 <http://maternalmortalitydata.org/mme.html>

COMMONLY USED APPROACHES TO ESTIMATING THE IMPACT OF CONTRACEPTIVE USE

	Adding It Up Guttmacher Institute	Impact 2 MSI	ImpactNow Futures Group	Reality Check EngenderHealth	FamPlan/LiST Futures Institute
PURPOSES	Illustrate the health and financial costs and benefits of modern contraceptive use in terms of outcomes averted through current or full (i.e. no unmet need) contraceptive use.	Estimate the current and potential future impacts of FP service provision on adverse outcomes averted through comparing current contraceptive use (modelled from service statistics) with no use; Estimate the quantity of contraceptive services needed to reach a specified contraceptive prevalence goal.	Estimate the health and economic impacts of contraceptive use in the near-term; automatic scenario comparisons facilitate creation of advocacy messages. Can also estimate impacts for a given FP budget.	Estimate the resources required to maintain or reach a specified contraceptive prevalence, increase by a certain annual rate, or meet unmet need; Estimate the potential impact of achieving this goal on averting adverse outcomes.	Estimate the resources required to meet contraceptive goals, such as reducing unmet need, achieving a specified modern contraceptive prevalence goal, or achieving a goal of reduced total fertility.
DATA INPUT REQUIREMENTS	No input required from user. Tabulations of most recent national surveys (e.g. DHS, MICS); estimations for missing data: – Method use; – Unmet need (by marital status and spacing/limiting); – Sub-regional estimates of unintended pregnancies; – Estimated proportions of pregnancy outcomes for birth, safe/unsafe abortion and miscarriage; – Maternal death rate and DALYs (for all women and those with abortions).	Required input from user: Quantity of services provided annually; or programme goal (CPR, additional users). The following inputs have defaults which can be overridden by the user: – Demographic data from UN Population Projections; – DHS and UN Population Division data on CPR, unmet need, % married; – Special studies for method effectiveness and discontinuation; – Costs & coverage by of maternal & neonatal interventions from UN Reproductive Health Costing Tool	The following inputs have default values, which can be overridden by the user: – Demographic data from UN Population Projections; – DHS and UN Population Division data on CPR, unmet need, % Married; – Special studies for method effectiveness and discontinuation; – Costs & coverage by of maternal & neonatal interventions from UN Reproductive Health Costing Tool and others; – Unit costs of FP provision (by method).	The following inputs have default values which can be overridden by the user: – Method discontinuation, CYP factors, per-unit costs, pregnancy rate, unintended pregnancy outcomes, maternal mortality ratio; – Tabulations of most recent national surveys (e.g. DHS, MICS) for CPR; – Demographic data from UN Population Projections.	The following inputs have default values preloaded, which cannot be overridden by the user: – Demographic data on population by age and sex in base year, TFR, age distribution of fertility, sex ratio at birth, life expectancy at birth, migration by age and sex from UN Population Projections. – DHS data on method mix, source mix, proximate determinants of fertility, costs of FP by method and source, method effectiveness, CYP factors, and percent of births that are high risk; – Data from special studies for method effectiveness and CYP factors.
OUTCOMES MEASURED	<i>Impacts estimated:</i> – Unintended pregnancies; – Unintended births (live & still); – Safe & unsafe abortions; – Miscarriages; – Maternal deaths; – DALYs; – Contraceptive service costs; – Maternal and newborn health care costs.	<i>Programme outcomes:</i> – FP users by method – Contribution to CPR – Services needed to reach goal <i>Impacts averted:</i> – Unintended pregnancies; – Live births; – Safe & unsafe abortions; – Maternal & child deaths; – DALYs; – Direct costs saved to families and healthcare systems (maternal & neonatal).	<i>Programme outcomes:</i> – FP users and acceptors, by method – CPR; – Unmet need; – FP budget required (depending on configuration chosen) <i>Impacts averted:</i> – Unintended pregnancies; – Live births; – Safe & unsafe abortions; – Maternal & child deaths; – DALYs; – Maternal & neonatal costs averted; – Cost-benefit ratio; – ICER (Incremental Cost-Effectiveness Ratio).	<i>Programme outcomes:</i> – FP users and acceptors; – Commodities Commodity costs <i>Impacts averted:</i> – Unintended pregnancies; – Live births; – Safe & unsafe abortions; – Maternal deaths.	<i>Programme outcomes:</i> – TFR; – CPR; – FP users and acceptors, by method and source; – Unmet need; – Average method effectiveness; – CYP; – Commodities required; – Contraceptive service costs; – Percentage of births with demographic risk factors; – Pregnancies; – Births by mother's age, birth interval and parity; – Abortions; – Maternal deaths, by type; – Infant and under 5 mortality rates <i>Impacts averted:</i> – Impacts can be calculated for any of the programme outcomes listed above
METHOD USED TO ESTIMATE NUMBER OF ABORTIONS AVERTED	Apply current proportion of sub-regional unintended pregnancies that end in abortions (safe & unsafe) to unintended pregnancies averted.	Apply sub-regional abortion ratios to the number of live births averted nationally; Live births averted calculated by subtracting estimated abortions, stillbirths, and miscarriages from pregnancies averted.	Apply sub-regional abortion ratios to the number of live births averted nationally; Live births averted calculated by subtracting estimated abortions, stillbirths, and miscarriages from pregnancies averted.	Apply current proportion of sub-regional unintended pregnancies that end in abortions (safe & unsafe) to unintended pregnancies averted.	Number of abortions calculated from an assumed total abortion rate, or as a percentage of unintended pregnancies.
METHOD USED TO ESTIMATE NUMBER OF MATERNAL DEATHS AVERTED	Apply mortality ratios for live births, safe & unsafe abortions averted; Ratios based on WHO country MMR, WHO regional unsafe abortion mortality and estimated safe abortion mortality.	Apply the national maternal mortality ratio to the number of live births averted Default MMR values from WHO provided.	Apply the national maternal mortality ratio to the number of live births averted. Default MMR values from WHO provided.	Apply MMR to number of live births averted. Default MMR values from WHO provided.	Apply MMR to the number of live births.
METHOD USED TO ESTIMATE NUMBER OF INFANT AND CHILD DEATHS AVERTED	Attribute infant mortality reduction to fewer births. Apply WHO or IHME mortality rates to unplanned births averted.	Country-specific coefficient applied to the number of live births; this coefficient estimates the effect of longer birth intervals on decreasing child mortality. Apply coefficient developed using PSI methodology for birth spacing impact.	Country-specific coefficient applied to the number of live births; this coefficient estimates the effect of longer birth intervals on decreasing child mortality. Apply coefficient developed using PSI methodology for birth spacing impact.	Infant and child deaths averted are not outputs of the tool.	Deaths averted result from two factors: 1) contraception leads to fewer births which result in fewer deaths, 2) contraception leads to fewer risky births (high parity, closely spaced, mother's age <18 or >35) which can reduce mortality rates and thus result in fewer deaths.

INFANT AND CHILD DEATHS AVERTED

The estimates of infant and child deaths averted are recognized as the weakest elements of these models because the relationship between contraceptive use and infant and child mortality is still being clarified. However, these impacts are requested by many users for advocacy purposes and so all models are looking to improve the estimation methods; for example, *Impact2* and *FamPlan* are exploring how contraceptive use changes the distribution of birth spacing risks and subsequent child mortality. *AIU* currently estimates infant deaths averted due to a reduction in the number of births (i.e. the demographic effect), rather than changes in risk and will explore integrating findings regarding changes in birth spacing.

IMPLICATIONS FOR MEASURING THE IMPACT OF CONTRACEPTIVE USE

These meetings and modelling exercises provided an important and unique opportunity for those involved in the development of estimation methodologies to come together and to discuss the various approaches used. This review process enabled the group to identify where and why differences existed and to seek alignment for harmonising the modelling approaches more closely. As a result, the estimation methodologies for each approach have been coordinated and consensus reached to ensure that, wherever possible, they are not in conflict. This alignment and consensus building process has strengthened the models by enabling their developers to benefit from each other's' experience and research. Moreover, decision-makers and managers using the different models can more clearly understand the assumptions behind each model so as to make informed choices between them.

MEETING PARTICIPANTS

JACOB ADETUNJI (USAID)
IAN ASKEW (Population Council)
ANGELA BASCHIERI (DFID)
ANN BIDDLECOM (UNPD)
JOHN BONGAARTS (Population Council)
WIN BROWN (Bill & Melinda Gates Foundation)
JACQUELINE E. DARROCH (Guttmacher Institute)
SENNEN HOUNTON (UNFPA)
ANRUDH JAIN (Population Council)
AMY RATCLIFFE (PSI)
MEGHAN REIDY (PSI)
AMANI SELIM (USAID)
SUSHEELA SINGH (Guttmacher Institute)
ELLEN SMITH (Futures Group)
JOHN STOVER (Futures Institute)
MICHELLE WEINBERGER (MSI)
MELANIE YAHNER (EngenderHealth)

STEP UP generates policy-relevant research to promote an evidence-based approach for improving access to family planning and safe abortion services.

We work in Bangladesh, northern India, Ghana, Kenya, and Senegal.

Population Council
Coordinating Partner

African Population and
Health Research Center
icddr,b

London School of Hygiene
and Tropical Medicine

Marie Stopes International
Partners in Population and
Development

Funded by