2016

Findings from landscape analysis in Bauchi on pre-eclampsia/ eclampsia

Population Council

Follow this and additional works at: https://knowledgecommons.popcouncil.org/departments_sbsr-rh

Part of the [Demography, Population, and Ecology Commons](#), [Family, Life Course, and Society Commons](#), [International Public Health Commons](#), and the [Maternal and Child Health Commons](#)

How does access to this work benefit you? Let us know!

Recommended Citation

"Findings from landscape analysis in Bauchi on pre-eclampsia/eclampsia," Ending Eclampsia Research Brief. Washington, DC: Population Council, 2016.

This Brief is brought to you for free and open access by the Population Council.

Despite global efforts to reduce preventable maternal and neonatal mortality, Nigeria's maternal mortality ratio is estimated at 576 deaths per 100,000 live births and neonatal death is estimated at 37 per 1,000 live births¹.

Maternal and newborn deaths due to pre-eclampsia and eclampsia (PE/E) are preventable, yet in Nigeria this is the most significant direct cause of maternal deaths.

To appreciate the enormity of this problem at country and state levels, a landscape analysis was conducted by the Population Council in 2015 on PE/E in seven states in Nigeria. The main objectives of the landscape analysis were:

- To understand the level of programmatic and policy support for PE/E prevention and treatment;
- To analyze the gaps in providers' knowledge and competence in preventing, detecting, and managing PE/E;
- To determine primary health care (PHC) facilities' capacities to manage PE/E;
- To assess community awareness, beliefs, and experiences around PE/E;
- To understand the volume of research on PE/E in the last 15 years; and
- To determine priority areas for research and programmatic interventions around PE/E.

The Ending Eclampsia project seeks to expand access to proven, underutilized interventions and commodities for the prevention, early detection, and treatment of pre-eclampsia and eclampsia and strengthen global partnerships.

PE/E IN BRIEF

- Pre-eclampsia is a condition in pregnant women marked by an increase in blood pressure and protein in urine after 20 weeks gestation.
- Providing high quality antenatal care improves the prevention and early detection of pre-eclampsia and can prevent its progression to eclampsia.
- Eclampsia is a life-threatening condition characterized by convulsions in women with PE.
- Women in developing countries are 300 times more likely to die from eclampsia than women in developed countries.
- Prescribing low-dose aspirin and calcium to at-risk women can prevent pre-eclampsia and eclampsia.
- Pre-eclampsia and eclampsia can be managed by administering anti-hypertensive drugs and magnesium sulphate ($MgSO_4$).
- $MgSO_4$ is the safest and most effective treatment for severe PE/E, and is one of 13 UN Life-Saving Commodities for Women and Children.
- PE/E and other hypertensive disorders in pregnancy increase the risk of pre-term births, which can lead to low birth weight, anemia, and stunting.
- Improved prevention, increased detection, and effective treatment of PE/E can prevent unnecessary maternal and newborn deaths.

FACILITY CAPACITY AND PREPAREDNESS

To assess institutional preparedness, researchers visited fourteen facilities in Bauchi State and recorded that only one (10%) of the facilities had guidelines available for management of pre-eclampsia, eight (57%) had all ANC equipment for the detection of PE/E and five (38%) use $MgSO_4$ for the treatment of eclampsia.

During these facility assessments, researchers determined whether the facilities had the key ANC equipment required to detect pre-eclampsia, manage severe PE/E, and monitor for $MgSO_4$ toxicity (figure 1).

FIGURE 1 Proportion of facilities with key equipment to detect PE and manage severe PE/E

Eight (57%) facility managers reported always using $MgSO_4$ to treat PE/E, xx (24%) said it is sometimes used, and xx (19%) reported that it is never used. When asked how they obtain $MgSO_4$, 18% of managers said they receive it routinely from central supply and 29% purchase locally, 47% ask clients to purchase it and 6% said they procure it elsewhere (such as NGOs).

Although hospitals have the capacity to provide signal functions for emergency obstetric and neonatal care (EmONC), Table 1 shows that PHCs have limited capacity for conducting even basic EmONC.

TABLE 1 Signal functions for emergency obstetric and newborn care (EmONC)

	Tertiary/Secondary health facilities (n=3)		Primary Health facilities (n=18)	
	Capacity	Done in the last three months	Capacity	Done in the last three months
Parenteral antibiotics	3	3	16	16
Parenteral oxytocics	3	3	16	16
Parenteral anticonvulsants ($MgSO_4$)	3	3	13	13
Manual removal of placenta	3	3	15	15
Removal of retained products	3	3	12	12
Assisted vaginal delivery	2	2	7	7
Blood transfusion	3	3	9	8

“It [national protocol] must be widely distributed so people will become aware, they get it, touch it, read it, it should be translated into various languages because of the many tribes in Nigeria, so they understand, and for it to reach a wider set of people.”

—POLICYMAKER, FOMWAN, BAUCHI

PROVIDER KNOWLEDGE AND SKILLS

Providers had good knowledge on signs and symptoms of PE (64%) and E (82%), but only 34.5% could correctly identify signs and symptoms of chronic hypertension and 6% could identify severe PE correctly.

FIGURE 2 Provider knowledge of drugs for PE/E prevention and management (n=55)

*Number of providers who had reported awareness of prophylactic drugs for PE/E

Researchers also assessed health providers' knowledge of drugs used for preventing and managing PE/E as well as calcium gluconate to treat $MgSO_4$ toxicity (figure 2).

Figure 2 shows limited provider knowledge on prophylactic use of calcium and aspirin for women at risk of PE and which anti-hypertensives (aldomet or nifedipine) to use for managing high BP during pregnancy. The Pritchard regimen for $MgSO_4$ administration is considered the 'gold standard' for

preventing and treating convulsions in severe PE/E, but few providers (7%) could accurately describe the appropriate doses of MgSO₄ or name the antidote for MgSO₄ toxicity, calcium gluconate (xx%).

QUALITY OF CARE

Quality of care was assessed through observations of 28

FIGURE 3 Percent of clients who received all eight ANC components

client-provider interactions and 28 client exit interviews with the same pregnant women. Researchers assessed eight essential components of quality antenatal care (ANC). Figure 3 describes the components that clients received.

In addition to the eight essential components of ANC, questions and tests should be conducted to assess a woman's risk of developing PE/E, to detect PE/E, and inform clients of the signs of impending eclampsia.

During 11% of the observed ANC consultations, providers performed the necessary checks to detect women at risk of developing PE. These included history of high blood pressure (BP) and diabetes, date of last delivery, client's parity, age and weight, edema of face, hands, legs, and ankles. In addition, 82% providers observed in Bauchi measured women's BP and checked urine for protein.

Only 32% of the providers advised the clients on the symptoms of impending eclampsia (severe headache, blurred vision, and pre-eclampsia with generalized body swelling).

COMMUNITY KNOWLEDGE AND PERCEPTIONS

The study also included in-depth interviews (IDIs) and focus group discussions (FGDs) with PE/E survivors,

community stakeholders, and families affected by PE/E.

Married men reported consuming concoctions, thinking, stress, an evil curse and family history of PE/E as causes of high blood pressure, swelling, and bleeding in pregnancy. The men also said that to prevent these complications, they should encourage their wives to exercise regularly, attend ANC and to take their medications correctly.

Women stated that high blood pressure during pregnancy is caused by headaches and dizziness. Women also discussed taking traditional medications at home, but that the condition wasn't cured until they went to the hospital.

“Stress causes high blood pressure, a pregnant mother needs to be examined alongside her unborn baby. High BP might be hereditary and it's when [BP] is high that it results in eclampsia.”

—MALE FGD PARTICIPANT, BAUCHI

Overall, the qualitative findings show that the signs and symptoms associated with PE/E are often attributed to other causes and community members often seek traditional or spiritual healing before medical care. Misconceptions, myths, and mistrust between communities and health providers negatively influence care-seeking behaviors.

SURVIVORS' EXPERIENCES

Interviews with survivors documented their care-seeking pathways, including their PE/E experience, availability and accessibility of essential services and commodities, and the outcomes of the pregnancy for mother and child. Survivors' experiences provide insight informing strategies to work more closely with communities and health facilities improve access to, and use of, quality care.

The women interviewed were similar in age, ages at marriage and first pregnancy, and education. The following quotation demonstrates lack of awareness and poor health seeking behavior in a young woman during her first pregnancy.

“My husband and I went to the hospital when my pregnancy was four months. At seven months, I returned for another antenatal care visit, they told me that my blood pressure had gone up to 140/120, I needed to take very good care of myself. At nine months, I started feeling pains in my legs. Upon reaching the hospital, I was told I still have some time. Unexpectedly, I felt a severe headache and my eyes felt heavy; I became unconscious. The baby died.”

—PE/E SURVIVOR, BAUCHI

DISCUSSION & CONCLUSION

This landscape analysis identified the gaps in facilities’ and providers’ capacities for preventing, detecting, and managing PE/E; it assessed community awareness, beliefs, and experiences of PE/E; and it determined the gaps and priority areas for research and programs to improve access to prevention and treatment.

It is clear, at the state policy level, that there is limited knowledge of what has been developed nationally; there is confusion about how policies, guidelines, and protocols reach not only the states but also the facilities themselves, where they don’t seem to exist at all.

For effective PE/E management, health facilities need to stock necessary drugs and ANC equipment, and institutionalize guidelines for PE/E treatment. Researchers noted irregular supply of basic drugs, equipment, and supplies, particularly MgSO_4 , patella hammers and the MgSO_4 antidote, calcium gluconate, which no facilities stocked.

Providers’ knowledge is generally poor in Bauchi; health care providers need to be trained and re-trained on PE prevention—including when to administer prophylactic drugs such as calcium supplements and aspirin—and its early detection, and how and when to administer anti-hypertensives and MgSO_4 for early treatment of the condition.

In addition to ensuring providers are adequately trained to administer MgSO_4 at the right time, and with the

proper doses, they also need to know the warning signs for MgSO_4 toxicity and its antidote, calcium gluconate. There is also limited capacity for newborn resuscitation.

The final, essential component to reduce mortality from PE/E is community awareness. Community members need to know the signs of PE/E and understand the danger it poses for mothers and babies so they can seek medical care promptly.

A multi-pronged approach is required to address the systems issues that create barriers to women’s access to health care during pregnancy. Detecting and managing PE/E early can improve the survival rate in women and babies in Nigeria.

RECOMMENDATIONS

- Advocacy for streamlining state procurement and link to a national or central distribution system;
- Connecting with United Nations Commission on Life-Saving Commodities (UNCoLSC) for Women and Children, especially for maternal drugs which include MgSO_4 ;
- Training on quality ANC: what, why, how, when and where;
- Introduce task-shifting policy focusing on MNH “policy implementation plan” to include anti-hypertensives and MgSO_4 ;
- Train PHC workers to detect and provide loading dose of MgSO_4 ;
- Community awareness campaigns on the importance of ANC and seeking care early; and
- Engage men to encourage their support of their wives.

RESOURCES

1. National Population Commission (NPC) [Nigeria] and ICF International. 2014. Nigeria Demographic and Health Survey 2013. Abuja, Nigeria and Rockville, Maryland, USA: NPC and ICF International.
2. Oladapo OT, Adetoro OO, Ekele BA, Chama C, Etuk SJ, Aboyeji AP, Onah HE, Abasiattai AM, Adamu AN, Adegbola O, Adeniran AS, Aimakhu CO, Akinsanya O, Aliyu LD, Ande AB, Ashimi A, Bwala M, Fabamwo A, Geidam AD, Ikechebelu JI, Imaralu JO, Kuti O, Nwachukwu D, Omo-Aghoja L, Tunau K, Tukur J, Umeora OUJ, Umezulike AC, Dada OA, Tunçalp O, Vogel JP, Gülmezoglu AM, Nigeria Near-miss and Maternal Death Surveillance Network. When getting there is not enough: a nationwide cross-sectional study of 998 maternal deaths and 1451 near-misses in public tertiary hospitals in a low-income country. BJOG 2015; DOI: 10.1111/1471-0528.13450.