


2016

# Findings from landscape analysis in Cross River on pre-eclampsia/eclampsia

Population Council

Follow this and additional works at: [https://knowledgecommons.popcouncil.org/departments\\_sbsr-rh](https://knowledgecommons.popcouncil.org/departments_sbsr-rh)

 Part of the [Demography, Population, and Ecology Commons](#), [Family, Life Course, and Society Commons](#), [International Public Health Commons](#), and the [Maternal and Child Health Commons](#)

---

## Recommended Citation

"Findings from landscape analysis in Cross River on pre-eclampsia/eclampsia," Ending Eclampsia Research Brief. Washington, DC: Population Council, 2016.

This Brief is brought to you for free and open access by the Population Council.

Despite global efforts to reduce preventable maternal and neonatal mortality, Nigeria's maternal mortality ratio is estimated at 576 deaths per 100,000 live births and neonatal death is estimated at 37 per 1,000 live births<sup>1</sup>.

Maternal and newborn deaths due to pre-eclampsia and eclampsia (PE/E) are preventable, yet in Nigeria this is the most significant direct cause of maternal deaths.

To appreciate the enormity of this problem at country and state levels, a landscape analysis was conducted by the Population Council in 2015 on PE/E in seven states in Nigeria. The main objectives of the landscape analysis were:

- To understand the level of programmatic and policy support for PE/E prevention and treatment;
- To analyze the gaps in providers' knowledge and competence in preventing, detecting, and managing PE/E;
- To determine primary health care (PHC) facilities' capacities to manage PE/E;
- To assess community awareness, beliefs, and experiences around PE/E;
- To understand the volume of research on PE/E in the last 15 years; and
- To determine priority areas for research and programmatic interventions around PE/E.

*The Ending Eclampsia project seeks to expand access to proven, underutilized interventions and commodities for the prevention, early detection, and treatment of pre-eclampsia and eclampsia and strengthen global partnerships.*

## PE/E IN BRIEF

- Pre-eclampsia is a condition in pregnant women marked by an increase in blood pressure and protein in urine after 20 weeks gestation.
- Providing high quality antenatal care improves the prevention and early detection of pre-eclampsia and can prevent its progression to eclampsia.
- Eclampsia is a life-threatening condition characterized by convulsions in women with PE.
- Women in developing countries are 300 times more likely to die from eclampsia than women in developed countries.
- Prescribing low-dose aspirin and calcium to at-risk women can prevent pre-eclampsia and eclampsia.
- Pre-eclampsia and eclampsia can be managed by administering anti-hypertensive drugs and magnesium sulphate (MgSO<sub>4</sub>).
- MgSO<sub>4</sub> is the safest and most effective treatment for severe PE/E, and is one of 13 UN Life-Saving Commodities for Women and Children.
- PE/E and other hypertensive disorders in pregnancy increase the risk of pre-term births, which can lead to low birth weight, anemia, and stunting.
- Improved prevention, increased detection, and effective treatment of PE/E can prevent unnecessary maternal and newborn deaths.

## FACILITY CAPACITY & PREPAREDNESS


To assess institutional preparedness, researchers visited 11 facilities in Cross River State and recorded that four (36%) of the facilities had guidelines available for management of pre-eclampsia, two (18%) had all ANC equipment for the detection of PE/E and five (46%) use MgSO<sub>4</sub> for the treatment of eclampsia.

“QUOTATION. QUOTATION.  
QUOTATION. QUOTATION.  
QUOTATION. QUOTATION.  
QUOTATION. QUOTATION.  
QUOTATION.”

—POLICYMAKER, CROSS RIVER

During these facility assessments, researchers determined whether the facilities had the key ANC equipment required to detect pre-eclampsia, manage severe pre-eclampsia and eclampsia, and monitor for MgSO<sub>4</sub> toxicity (figure 1).

**FIGURE 1** Proportion of facilities with key equipment to detect PE and manage severe PE/E




Four (36%) facility managers reported always using MgSO<sub>4</sub> to treat pre-eclampsia and eclampsia, one (9%) said it is sometimes used, and six (55%) reported that it is never used. When asked about how they obtain MgSO<sub>4</sub> supplies at the facility, the Council found that 80% receive the drug as part of the regular central supply and 20% require their clients to purchase it from the market.

## PROVIDER KNOWLEDGE & SKILLS

Fifty-nine healthcare providers from Cross River participated in the study and answered questions about different aspects of PE/E prevention, detection, and treatment. Of the providers interviewed, 50.8% could correctly identify chronic hypertension in pregnancy, while 64% correctly identified signs and symptoms of pre-eclampsia, yet only 5% could identify severe pre-eclampsia, and 71% could correctly identify the signs and symptoms of eclampsia.

Researchers also assessed health care providers' knowledge of drugs used for preventing and managing PE/E as well as calcium gluconate to treat MgSO<sub>4</sub> toxicity (figure 2).

**FIGURE 2** Provider knowledge of drugs for PE/E prevention and management (n=59)



\*Number of providers who had reported awareness of prophylactic drugs for PE/E

It is clear from the results, shown in figure 2, that providers are unaware of the prophylactic use of calcium and aspirin for women at risk of pre-eclampsia: only two (18%) knew that anti-hypertensives (aldomet or nifedipine) can be used to manage high blood pressure in pregnant women.

The Pritchard regimen for MgSO<sub>4</sub> administration is considered the 'gold standard' for preventing and treating convulsions in severe PE/E, but only a small percentage of providers (10%) could accurately describe the appropriate loading and maintenance doses of MgSO<sub>4</sub>.

Meanwhile, only three percent of providers could name the appropriate antidote for MgSO<sub>4</sub> toxicity: calcium gluconate.



are cheap and obtainable, are often available in maternity emergency trays at most facilities and, if not, can be purchased externally. It is imperative that future interventions targeting providers, especially at primary and secondary facilities, include training on anti-hypertensives for pregnant women with PE/E.

For effective PE/E management, health facilities need to stock the necessary drugs and ANC equipment, and institutionalize guidelines for PE/E treatment. Providers need training and re-training on detecting PE/E, and how and when to administer appropriate drugs (prophylactics, anti-hypertensives, MgSO<sub>4</sub>) to prevent and manage it. With few providers understanding these elements of PE/E detection and treatment, it is not surprising these pregnancy complications account for more maternal deaths in Nigeria than any other direct cause, including postpartum hemorrhage<sup>2</sup>.

In addition to ensuring that health care providers are adequately trained to administer MgSO<sub>4</sub> at the right time and with the proper doses, they also need to know the warning signs for MgSO<sub>4</sub> toxicity and its antidote, calcium gluconate.

The final, essential component to reduce mortality from PE/E is community awareness. Community members need to know the signs of PE/E and understand the danger it poses for mothers and babies so they can seek medical care promptly.

## CONCLUSION

Maternal and newborn deaths due to PE/E are preventable: by increasing community awareness of the condition, improving antenatal care quality, and scaling up proven best practices to prevent pre-eclampsia's escalation to severe pre-eclampsia and eclampsia. By detecting and managing pre-eclampsia, Ending Eclampsia can improve the survival rate of women and babies in Nigeria and other developing countries.

## RESOURCES

1. National Population Commission, Federal Republic of Nigeria, and ICF International. (2014). Nigeria Demographic and Health Survey 2013.
2. Oladapo, O., Adetoro, O., et al. (2015). When getting there is not enough: a nationwide cross-sectional study of 998 maternal deaths and 1451 near-misses in public tertiary hospitals in a low-income country. *BJOG: An International Journal of Obstetrics & Gynaecology*.